

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

COPEIA 15

No. 6. M. femur-rubrum, 5; Ipidae (Bark Beetle) 1; Wasp, 1.

No. 7. M. femur-rubrum, 1; L. 10-lineata, 6.

No. 8. M. femur-rubrum, 2; L. 10-lineata, 1.

No. 9. L. 10-lineata, 2; Gryllus assimilis, 1; Nemobius fasciatus, 2.

No. 10. M. femur-rubrum, 3; L. 10-lineata, 1.

No. 11. M. femur-rubrum, 1; L. 10-lineata, 2.

No. 12. M. femur-rubrum, 5.

No. 13. M. femur-rubrum, 4; Dendroctonus sp? (Bark Beetle) 1.

No. 14. M. femur-rubrum, 5; L. 10-lineata, 1.

No. 15. M. femur-rubrum, 3; L. 10-lineata, 1.

No. 16. M. femur-rubrum, 4.

No. 17. M. femur-rubrum, 5; L. 10-lineata, 2.

No. 18. Gryllus assimilis, 1; L. 10-lineata, 1; Agalena naevia, 2.

No. 19. M. femur-rubrum, 5.

No. 20. M. femur-rubrum, 8; L. 10-lineata, 1; Harpalus caliginosus (Ground Beetle), 1.

No. 21. M. femur-rubrum, 2; L. 10-lineata, 1.

No. 22. M. femur-rubrum. 3.

No. 23. M. femur-rubrum, 3; Mamestra assimilis larva (Cutworm) 1; Gryllus assimilis, 1; Euschistus variolarius (Bug) 1; Pentotomid nymph, 1.

No. 24. L. 10-lineata, 4.

No. 25. M. femur-rubrum, 3; Mamestra assimilis (Cutworm) 3; Agalena naevia, 1.

A. Brooker Klugh, Queen's University, Canada.

EGG-LAYING BY THE "HORNED TOAD", PHRYNOSOMA CORNUTUM.

According to Gadow (Cambridge Natural History, p. 533 of vol. Amphibia and Reptiles) all the species of *Phrynosoma* or Horned Lizards are viviparous, in which they differ from nearly all of

16 COPEIA

the other Iguanidae. A female specimen of the Texas "Horned Toad," P. cornutum, whose previous history was unknown, was given the writer late in the spring of 1921. During the night of June 7 to 8 this animal laid three eggs, whose yellowish-white, leathery shells measured 10 mm. x 16 mm. in size.

On the evening of June 11 the animal was found dead in its cage; it had probably been dead for some hours. On opening the body it was found to contain 34 eggs, similar to the three that had been laid four days before.

The writer has made no observations upon the breeding habits of these lizards, and he would like to know whether they commonly vary their sup-

posed viviparous habits in this way.*

Since no embryos were found in any of a number of the eggs that were examined, it is likely that fertilization had not taken place. In this event the eggs would not, of course, develop and they would have to be laid in order to be gotten from the body.

It is possible that the animal had been in captivity for some time before coming into the writer's

hands, and had thus escaped impregnation.

The bulk of the eggs was so enormous in comparison to the size of the animal that their resorption would seem impossible.

ALBERT M. REESE. West Virginia University.

^{* [}Our most detailed account of the egg laying of P. cornutum is that of Strecker (1908, Proc. Biol. Soc. Wash., XXI, pp. 165-169). Editors.1